



National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet
Search PubMed	for #19 AND #20 AND #21				Go Clear
<input checked="" type="checkbox"/> Limits		Index	History	Clipboard	

Entrez PubMed

PubMed Services

Related Resources

- The Clipboard will hold a maximum of 500 items
- Clipboard items will be lost after one hour of inactivity.

Display	Summary	<input checked="" type="checkbox"/> Save	Text	Order	Remove from Clipboard
Show: 20	Items 1-16 of 16				One page.

- ☐ 1 : Youngs SJ, Ali SA, Taub DD, Rees RC. Related Articles
Chemokines induce migrational responses in human breast carcinoma cell lines.
Int J Cancer. 1997 Apr 10;71(2):257-66.
PMID: 9139852; UI: 97284438
- ☐ 2 : Zhang L, Khayat A, Cheng H, Graves DT. Related Articles
The pattern of monocyte recruitment in tumors is modulated by MCP-1 expression and influences the rate of tumor growth.
Lab Invest. 1997 Apr;76(4):579-90.
PMID: 9111518; UI: 97265612
- ☐ 3 : Kurtz RM, Elner VM, Bian ZM, Strieter RM, Kunkel SL, Elner SG. Related Articles
Dexamethasone and cyclosporin A modulation of human retinal pigment epithelial cell monocyte chemotactic protein-1 and interleukin-8.
Invest Ophthalmol Vis Sci. 1997 Feb;38(2):436-45.
PMID: 9040477; UI: 97192885
- ☐ 4 : Van Snick J, Houssiau F, Proost P, Van Damme J, Renauld JC. Related Articles
I-309/T cell activation gene-3 chemokine protects murine T cell lymphomas against dexamethasone-induced apoptosis.
J Immunol. 1996 Sep 15;157(6):2570-6.
PMID: 8805659; UI: 96399088
- ☐ 5 : Jordan NJ, Watson ML, Westwick J. Related Articles
The protein phosphatase inhibitor calyculin A stimulates chemokine production by human synovial cells.
Biochem J. 1995 Oct 1;311 (Pt 1):89-95.
PMID: 7575485; UI: 96003901
- ☐ 6 : Liebler JM, Kunkel SL, Burdick MD, Standiford TJ, Rolfe MW, Strieter RM. Related Articles
Production of IL-8 and monocyte chemotactic peptide-1 by peripheral blood monocytes. Disparate responses to phytohemagglutinin and lipopolysaccharide.
J Immunol. 1994 Jan 1;152(1):241-9.
PMID: 8254194; UI: 94075828
- ☐ 7 : Antony VB, Godbey SW, Kunkel SL, Hott JW, Hartman DL, Burdick MD, Strieter RM. Related Articles
Recruitment of inflammatory cells to the pleural space. Chemotactic cytokines, IL-8,

Art Unit: 1644

characterization of the polypeptide encoded thereby by employing those methods that were old and well known in the art at of molecular biology at the time the invention was made would have been prima facie obvious to an artisan having ordinary skill. The motivation is to further characterize the properties of the isolated channel protein in a lipid bilayer system.

Conclusion

8. No claims are allowed.

Advisory Information

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Joseph F. Murphy whose telephone number is 703-305-7245. The examiner can normally be reached on M-F 8:30-5:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Christina Chan can be reached on 703-308-3973. The fax phone numbers for the organization where this application or proceeding is assigned are 703-305-3014 for regular communications and 703-308-0294 for After Final communications.

Any inquiry of a general nature or relating to the status of this application or proceeding should be directed to the receptionist whose telephone number is 703-308-0196.

Joseph F. Murphy, Ph. D.
Patent Examiner
Art Unit 1644
December 27, 1999

and monocyte chemotactic peptide-1 in human pleural fluids.
J Immunol. 1993 Dec 15;151(12):7216-23.
PMID: 8258721; UI: 94081372

- ☐ 8 : Rowell DL, Eckmann L, Dwinell MB, Carpenter SP, Raucy JL, Yang SK, Kagnoff MF. Related Articles
Human hepatocytes express an array of proinflammatory cytokines after agonist stimulation or bacterial invasion.
Am J Physiol. 1997 Aug;273(2 Pt 1):G322-32.
PMID: 9277410; UI: 97423468
- ☐ 9 : Gautam SC, Noth CJ, Janakiraman N, Pindolia KR, Chapman RA. Related Articles
Induction of chemokine mRNA in bone marrow stromal cells: modulation by TGF-beta 1 and IL-4.
Exp Hematol. 1995 Jun;23(6):482-91.
PMID: 7768303; UI: 95285906
- ☐ 10 : Casselmann BL, Kilgore KS, Miller BF, Warren JS. Related Articles
Antibodies to neutrophil cytoplasmic antigens induce monocyte chemoattractant protein-1 secretion from human monocytes.
J Lab Clin Med. 1995 Nov;126(5):495-502.
PMID: 7595035; UI: 96074349
- ☐ 11 : Tanaka H, Minoshima T, Endo N. Related Articles
The effect of a synthetic 7-thiaprostaglandin E1 derivative, TEI-6122, on monocyte chemoattractant protein-1 induced chemotaxis in THP-1 cells.
Br J Pharmacol. 1995 Oct;116(4):2298-302.
PMID: 8564263; UI: 96120619
- ☐ 12 : Wenzel UO, Fouqueray B, Grandaliano G, Kim YS, Karamitsos C, Valente AJ, Abboud HE. Related Articles
Thrombin regulates expression of monocyte chemoattractant protein-1 in vascular smooth muscle cells.
Circ Res. 1995 Sep;77(3):503-9.
PMID: 7641321; UI: 95368769
- ☐ 13 : Gong JH, Clark-Lewis I. Related Articles
Antagonists of monocyte chemoattractant protein 1 identified by modification of functionally critical NH2-terminal residues.
J Exp Med. 1995 Feb 1;181(2):631-40.
PMID: 7836918; UI: 95138695
- ☐ 14 : Zhang YJ, Rutledge BJ, Rollins BJ. Related Articles, Protein
Structure/activity analysis of human monocyte chemoattractant protein-1 (MCP-1) by mutagenesis. Identification of a mutated protein that inhibits MCP-1-mediated monocyte chemotaxis.
J Biol Chem. 1994 Jun 3;269(22):15918-24.
PMID: 8195247; UI: 94253189
- ☐ 15 : Shyy YJ, Li YS, Kolattukudy PE. Related Articles, Protein, Nucleotide
Structure of human monocyte chemotactic protein gene and its regulation by TPA.
Biochem Biophys Res Commun. 1990 Jun 15;169(2):346-51.
PMID: 2357211; UI: 90290466

Art Unit: 1644

vector pT7T3. The polypeptide encoded by this polynucleotide would be identical to a fragment of the polypeptide set forth in SEQ ID NO: 2, thus anticipating claims 5-8.

b. Claims 2, 5-8 rejected under 35 U.S.C. 102(b) as being anticipated by FR 2744730-A1 (1997).

FR 2744730-A1 discloses the cloning of a cDNA sequence encoding a TWIK1 potassium channel. The TWIK1 channel protein disclosed in FR 2744730-A1 contains a stretch of bases from 529-549 that is identical to the polynucleotide sequence set forth in SEQ ID NO: 1 (see Sequence Comparison B, underlined region, attached), thus anticipating claim 2(i). Furthermore, the polynucleotide disclosed in FR 2744730-A1 was cloned into an expression vector, and expressed in oocytes (page 20, first paragraph), thus anticipating claims 5-8.

Claim Rejections - 35 USC § 103

7. The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negated by the manner in which the invention was made.

a. Claims 2 and 5-9 are rejected under 35 U.S.C. 103(a) as being unpatentable over FR 2744730-A1 (1997).

The disclosure of FR 2744730-A1 is set forth above (section 6b.). However, FR 2744730-A1 does not teach a method for producing the polypeptide encoded by the disclosed polynucleotide. To have incorporated the recombinant DNA identified in the disclosure of FR 2744730-A1 into an expression vector and host cell to facilitate the production and

☐ 16 : Yoshimura T, Yuhki N, Moore SK, Appella E, Lerman MI, Leonard EJ. Related Articles, Protein, Nucleotide

Human monocyte chemoattractant protein-1 (MCP-1). Full-length cDNA cloning, expression in mitogen-stimulated blood mononuclear leukocytes, and sequence similarity to mouse competence gene JE.
FEBS Lett. 1989 Feb 27;244(2):487-93.
PMID: 2465924; UI: 89153605

Display	Summary	<input type="checkbox"/>	Save	Text	Order	Remove from Clipboard
Show: 20	<input type="checkbox"/>	Items 1-16 of 16				One page.

Revised: January 10, 2000.

Disclaimer | Write to the Help Desk
NCBI | NLM | NIH

Art Unit: 1644

under conditions of moderate stringency, for example, would fail to hybridize under conditions of high stringency. The metes and bounds of the claim thus cannot be ascertained. This rejection could be obviated by supplying specific conditions supported by the specification which Applicant considers to be "stringent". Applicant should specifically point out the support for any amendments made to the disclosure. See MPEP 714.02 and 2163.06

Claim 2, sections (i) and (s), is indefinite in the recitation of the term "fragments". This language is vague and indefinite since it encompasses potentially any portion of the polypeptide including a single amino acid. There is no guidance provided as to what specific sequences the term "fragments" refers to. Therefore, the metes and bounds of the claim are unclear.

Claim Rejections - 35 USC § 102

6. The following is a quotation of the appropriate paragraphs of 35 U.S.C. 102 that form the basis for the rejections under this section made in this Office action:

A person shall be entitled to a patent unless –

(b) the invention was patented or described in a printed publication in this or a foreign country or in public use or on sale in this country, more than one year prior to the date of application for patent in the United States.

a. Claims 2 and 5-8 are rejected under 35 U.S.C. 102(b) as being anticipated by Marra et al (1996).

Marra et al. discloses the cloning of a cDNA which is identical from bases 309 to 331 to the sequence set forth in SEQ ID NO: 1 (see Sequence Comparison A, underlined region, attached). Since this stretch of bases is more than 15 bases long, it is within the limitations of claim 2(i). Furthermore, the sequence disclosed in Marra et al. was cloned into the expression